



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 4, 2026 – 08:56 AM EDT

PDB ID : 2ODE / pdb_00002ode
Title : Crystal structure of the heterodimeric complex of human RGS8 and activated Gi alpha 3
Authors : Gileadi, C.; Soundararajan, M.; Turnbull, A.P.; Elkins, J.M.; Papagrigoriou, E.; Pike, A.C.W.; Bunkoczi, G.; Gorrec, F.; Umeano, C.; von Delft, F.; Weigelt, J.; Edwards, A.; Arrowsmith, C.H.; Sundstrom, M.; Doyle, D.A.; Structural Genomics Consortium (SGC)
Deposited on : 2006-12-22
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	350	
1	C	350	
2	B	141	

Continued on next page...

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
 Validation Pipeline (wwPDB-VP) : 2.50

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	141	3% 75% 6% 18%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7749 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Guanine nucleotide-binding protein G(k) subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	316	Total	C	N	O	S	0	2	0
			2508	1597	416	479	16			
1	C	317	Total	C	N	O	S	0	4	0
			2511	1599	417	478	17			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	SER	-	expression tag	UNP P08754
A	3	MET	-	expression tag	UNP P08754
A	351	SER	-	expression tag	UNP P08754
C	2	SER	-	expression tag	UNP P08754
C	3	MET	-	expression tag	UNP P08754
C	351	SER	-	expression tag	UNP P08754


- Molecule 2 is a protein called Regulator of G-protein signaling 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	131	Total	C	N	O	S	0	3	0
			1064	677	175	208	4			
2	D	116	Total	C	N	O	S	0	1	0
			943	607	154	178	4			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	40	SER	-	expression tag	UNP P57771
B	41	MET	-	expression tag	UNP P57771
D	40	SER	-	expression tag	UNP P57771
D	41	MET	-	expression tag	UNP P57771

- Molecule 3 is TETRAFLUOROALUMINATE ION (CCD ID: ALF) (formula: AlF_4).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	Al	F	0	0
			5	1	4		
3	C	1	Total	Al	F	0	0
			5	1	4		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	C	1	Total	Mg	0	0
			1	1		

- Molecule 5 is GUANOSINE-5'-DIPHOSPHATE (CCD ID: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total 28	C 10	N 5	O 11	P 2	0	0
5	C	1	Total 28	C 10	N 5	O 11	P 2	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	261	Total O 261 261	0	0
6	B	99	Total O 99 99	0	0
6	C	213	Total O 213 213	0	0
6	D	82	Total O 82 82	0	0

- Molecule 1: Guanine nucleotide-binding protein G(k) subunit alpha



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	112.84Å 130.22Å 68.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.40 – 1.90 29.40 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (29.40-1.90) 99.9 (29.40-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.09 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.180 , 0.211 0.184 , 0.213	Depositor DCC
R_{free} test set	4021 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	30.5	Xtriage
Anisotropy	0.041	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 45.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7749	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.15% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GDP, ALF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	0/2559	0.85	0/3457
1	C	0.84	0/2568	0.85	0/3472
2	B	0.85	0/1096	0.85	1/1480 (0.1%)
2	D	0.87	1/967 (0.1%)	0.91	1/1302 (0.1%)
All	All	0.84	1/7190 (0.0%)	0.86	2/9711 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	106	HIS	C-O	-5.10	1.18	1.24

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	106	HIS	N-CA-C	5.35	117.11	111.28
2	D	128	ARG	NE-CZ-NH2	5.33	124.00	119.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	170	ASP	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2508	0	2443	10	0
1	C	2511	0	2425	16	0
2	B	1064	0	997	10	0
2	D	943	0	897	7	0
3	A	5	0	0	0	0
3	C	5	0	0	0	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	A	28	0	12	0	0
5	C	28	0	12	0	0
6	A	261	0	0	5	0
6	B	99	0	0	3	0
6	C	213	0	0	3	0
6	D	82	0	0	3	0
All	All	7749	0	6786	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:47:THR:HG22	2:B:175:SER:OG	1.44	1.16
2:D:106:HIS:HE1	6:D:226:HOH:O	1.61	0.82
2:B:47:THR:HG23	2:B:176:GLN:HG3	1.74	0.70
1:C:248:LYS:NZ	6:C:3202:HOH:O	2.24	0.70
2:B:106:HIS:HE1	6:B:268:HOH:O	1.77	0.68
2:B:52:ARG:HG3	2:B:59:VAL:HG11	1.82	0.62
2:D:84[A]:GLU:OE1	2:D:119:ARG:NH1	2.34	0.61
1:C:162[A]:ILE:HD12	1:C:167:TYR:CZ	2.37	0.60
1:C:74:TYR:HE1	1:C:162[A]:ILE:HG22	1.69	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:47:THR:HG22	2:B:175:SER:CB	2.34	0.56
1:C:330:LYS:O	1:C:334:PHE:HD2	1.88	0.56
2:B:60:LEU:C	2:B:60:LEU:HD23	2.31	0.56
1:A:347:ASN:CG	6:A:3058:HOH:O	2.49	0.54
2:D:168:TYR:O	2:D:171:LEU:HB2	2.08	0.54
1:C:143:SER:HA	1:C:146:TYR:CE1	2.43	0.53
1:A:297:GLU:HG2	6:A:3076:HOH:O	2.10	0.52
2:B:106:HIS:CE1	6:B:268:HOH:O	2.58	0.52
1:A:254:CYS:SG	1:A:319:ILE:HD11	2.51	0.51
2:D:106:HIS:CE1	6:D:226:HOH:O	2.46	0.51
2:B:101:LEU:C	2:B:101:LEU:HD23	2.34	0.51
1:A:38:LEU:CD2	1:A:50:VAL:HG23	2.42	0.49
2:B:47:THR:O	2:B:51:THR:HG23	2.12	0.49
1:A:60:GLY:CA	6:A:3062:HOH:O	2.60	0.48
1:A:143:SER:HA	1:A:146:TYR:CE1	2.48	0.48
1:C:226:ALA:HB1	1:C:270:LYS:HD2	1.97	0.46
1:C:110:LEU:HB2	1:C:123:LEU:HD13	1.97	0.45
1:C:162[A]:ILE:HD12	1:C:167:TYR:CE1	2.52	0.45
2:D:133:LYS:CE	6:D:252:HOH:O	2.65	0.44
1:C:38:LEU:HD23	1:C:50:VAL:HG23	2.00	0.44
1:A:60:GLY:HA2	6:A:3062:HOH:O	2.18	0.43
1:C:162[A]:ILE:HG23	1:C:167:TYR:CE2	2.53	0.43
2:D:101:LEU:C	2:D:101:LEU:HD23	2.44	0.43
1:A:89:GLY:N	6:A:3248:HOH:O	2.49	0.43
2:B:128:ARG:HD3	6:B:220:HOH:O	2.20	0.42
1:C:119:MET:HA	6:C:3163:HOH:O	2.18	0.42
1:A:71:VAL:HG12	1:A:117:GLY:CA	2.49	0.41
1:C:284:THR:HA	1:C:287:TYR:O	2.19	0.41
1:C:34:VAL:HG13	1:C:219:THR:HB	2.01	0.41
1:A:106:GLN:NE2	1:A:122:GLU:OE2	2.49	0.41
1:C:187:THR:HG22	6:C:3033:HOH:O	2.19	0.41
1:C:105:ARG:O	1:C:109:VAL:HG23	2.21	0.40
2:D:170:ASP:O	2:D:171:LEU:C	2.64	0.40
1:C:38:LEU:CD2	1:C:50:VAL:CG2	2.99	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	316/350 (90%)	312 (99%)	4 (1%)	0	100	100
1	C	319/350 (91%)	314 (98%)	5 (2%)	0	100	100
2	B	132/141 (94%)	130 (98%)	2 (2%)	0	100	100
2	D	115/141 (82%)	113 (98%)	2 (2%)	0	100	100
All	All	882/982 (90%)	869 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	268/308 (87%)	266 (99%)	2 (1%)	76	78
1	C	264/308 (86%)	259 (98%)	5 (2%)	50	47
2	B	112/128 (88%)	111 (99%)	1 (1%)	70	73
2	D	98/128 (77%)	97 (99%)	1 (1%)	68	70
All	All	742/872 (85%)	733 (99%)	9 (1%)	63	63

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	LEU
1	A	115	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	46	SER
1	C	38	LEU
1	C	105	ARG
1	C	115	GLU
1	C	158	ASP
1	C	347	ASN
2	D	115	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	79	GLN
2	B	116	GLN
1	C	171	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ALF	A	2001	6,5	4,4,4	1.28	0	-		
5	GDP	A	1001	4,3	29,30,30	1.19	3 (10%)	45,47,47	1.45	7 (15%)
3	ALF	C	2002	6,5	4,4,4	1.28	0	-		
5	GDP	C	1002	4,3	29,30,30	1.31	5 (17%)	45,47,47	1.53	8 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GDP	A	1001	4,3	-	1/16/32/32	0/3/3/3
5	GDP	C	1002	4,3	-	1/16/32/32	0/3/3/3

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	1002	GDP	C5-C4	3.01	1.47	1.38
5	A	1001	GDP	C5-C4	3.00	1.47	1.38
5	C	1002	GDP	C6-N1	-2.79	1.33	1.38
5	A	1001	GDP	C5-N7	-2.32	1.34	1.39
5	C	1002	GDP	C5-N7	-2.26	1.34	1.39
5	C	1002	GDP	C4-N9	-2.24	1.32	1.38
5	C	1002	GDP	C2-N1	-2.10	1.32	1.37
5	A	1001	GDP	PA-O3A	2.00	1.61	1.59

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	1002	GDP	C5-C4-N3	-4.34	121.49	128.39
5	C	1002	GDP	N9-C4-N3	4.09	134.12	125.95
5	A	1001	GDP	C5-C4-N3	-3.71	122.48	128.39
5	A	1001	GDP	N9-C4-N3	3.70	133.34	125.95
5	C	1002	GDP	C6-C5-N7	3.39	136.45	130.29
5	C	1002	GDP	C2-N3-C4	3.22	117.84	112.30
5	A	1001	GDP	O6-C6-C5	-2.94	118.77	126.53
5	A	1001	GDP	C6-C5-N7	2.59	135.00	130.29
5	A	1001	GDP	C2-N3-C4	2.55	116.70	112.30
5	C	1002	GDP	C6-C5-C4	-2.46	115.14	118.83
5	A	1001	GDP	C8-N9-C4	2.32	110.38	106.03
5	C	1002	GDP	C8-N9-C4	2.32	110.38	106.03
5	C	1002	GDP	O6-C6-C5	-2.30	120.47	126.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1001	GDP	O6-C6-N1	2.20	124.25	120.11
5	C	1002	GDP	C5-C6-N1	2.05	118.47	113.25

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1001	GDP	PA-O3A-PB-O3B
5	C	1002	GDP	PA-O3A-PB-O3B

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	316/350 (90%)	0.44	8 (2%) 58 62	21, 34, 42, 58	2 (0%)
1	C	317/350 (90%)	0.54	7 (2%) 62 66	20, 34, 43, 56	4 (1%)
2	B	131/141 (92%)	1.03	17 (12%) 7 7	21, 34, 40, 45	3 (2%)
2	D	116/141 (82%)	0.67	4 (3%) 48 51	20, 34, 40, 44	1 (0%)
All	All	880/982 (89%)	0.59	36 (4%) 41 44	20, 34, 43, 58	10 (1%)

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	65	TYR	4.0
1	A	112	GLY	3.9
2	D	171	LEU	3.7
2	B	125	PHE	3.7
2	B	123	ILE	3.4
2	B	127	THR	3.2
1	A	111	ALA	3.1
1	A	113	SER	2.9
1	C	118	VAL	2.8
1	A	348	LEU	2.8
2	B	115	VAL	2.8
1	A	118	VAL	2.7
1	C	111	ALA	2.7
2	B	122	ASN	2.7
1	A	59	ASP	2.6
2	D	169	LEU	2.6
1	C	117	GLY	2.6
2	B	113	VAL	2.6
2	D	168	TYR	2.5
2	B	46	SER	2.5
2	B	48	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	121	VAL	2.3
2	B	47	THR	2.3
2	B	128	ARG	2.3
2	D	150	VAL	2.3
2	B	124	ASP	2.3
2	B	176	GLN	2.2
1	C	96	GLY	2.2
1	C	31	ALA	2.2
1	C	168	ILE	2.2
1	A	117	GLY	2.1
2	B	130	ALA	2.1
2	B	126	GLN	2.1
1	A	315	ASP	2.1
2	B	131	THR	2.1
1	C	110	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ALF	C	2002	5/5	0.97	0.12	22,22,24,27	0
4	MG	C	3002	1/1	0.97	0.09	24,24,24,24	0
5	GDP	A	1001	28/28	0.97	0.08	17,21,24,25	0
5	GDP	C	1002	28/28	0.97	0.08	18,22,24,25	0
4	MG	A	3001	1/1	0.98	0.11	20,20,20,20	0
3	ALF	A	2001	5/5	0.98	0.11	20,21,22,23	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around GDP A 1001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around GDP C 1002:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.